

**(19) World Intellectual Property  
Organization  
International Bureau**



## PCT

**(10) International Publication Number**  
**WO 2004/044205 A1**

[KR/KR]; 13-1101 Hyundaiprime Apt., Guwui-dong, Gwangjin-gu, 143-761 Seoul (KR). **KIM, Mi-Jung** [KR/KR]; 467-7, Seongnae-1dong, Kangdong-gu, 134-031 Seoul (KR). **KIM, Hee-Ja** [KR/KR]; Rm.103 Daeyang Yeonrib 606-10, Shinwol2-dong, Yangcheon-gu, 158-092 Seoul (KR). **CHUNG, Chung-Han** [KR/KR]; Rm.111 Goejeong Jayu3cha Apt. 1065, Goejeong1-dong, Saha-gu, 604-081 Busan (KR). **PYEE, Jae-Ho** [KR/KR]; 1241-805 Mokryun Apt., Sanbon-dong, 435-040 Gunpo-city, Kyungki-do (KR). **HYUNG, Nam-In** [KR/KR]; 101-103 Hyundai Apt., Ssangyong-dong, 330-090 Cheonan-city, Chungcheongnam-do (KR).

(74) Agent: LEE, Young-Pil; The Cheonghwa Building, 1571-18, Seocho-dong Seocho-gu, 137-874 Seoul (KR).

**(81) Designated States (*national*):** JP, US.

**Published:**  
— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

R DERIVED FROM SESAME AND SEED-SPECIFIC EXPRES-

[illegible]

derived from sesame microsomal oleic acid desaturase (Si-FAD2) expression vector including the promoter and/or the intron, and a transgene. Therefore, a useful product can be produced in a seed-specific manner. Also, the promoter can be used together with the intron for the expression of an inserted gene in a seed. Therefore, it is very useful in developing a transgene in a seed-specific manner.

derived from sesame microsomal oleic acid desaturase (Si-FAD2) expression vector including the promoter and/or the intron, and a transgene. Therefore, a useful product can be produced in a seed-specific manner. Also, the promoter can be used together with the intron for the expression of an inserted gene in a seed. Therefore, it is very useful in developing a transgene in a seed-specific manner.

derived from sesame microsomal oleic acid desaturase (Si-FAD2) expression vector including the promoter and/or the intron, and a transgene. Therefore, a useful product can be produced in a seed-specific manner. Also, the promoter can be used together with the intron for the expression of an inserted gene in a seed. Therefore, it is very useful in developing a transgene in a seed-specific manner.

**(S7) Abstract:** Provided are a seed-specific expression promoter derived from sesame microsomal oleic acid desaturase (Si-FAD2) gene, an intron for expression enhancement, a seed-specific expression vector including the promoter and/or the intron, and a transgenic plant transformed with the seed-specific expression vector. Therefore, a useful product can be produced in a seed-specific manner or a common product in a seed can be functionally modified. Also, the promoter can be used together with the intron for expression enhancement, thereby increasing the expression level of an inserted gene in a seed. Therefore, it is very useful in development of a transgenic plant which induces large-scale expression of a foreign gene in a seed-specific manner.

WO 2004/044205

12/pfs  
10/534128  
JC06 Rec'd PCT/PTO 06 MAY 2005  
PCT/KR2003/002415

**PLANT SEED-SPECIFIC EXPRESSION PROMOTER DERIVED FROM  
SESAME AND SEED-SPECIFIC EXPRESSION VECTOR  
COMPRISING THE PROMOTER**

5 Technical Field

The present invention relates to a plant seed-specific expression promoter derived from a genomic gene (*Si-FAD2*) encoding the microsomal oleic acid desaturase of sesame and an intron for expression enhancement. The present invention also relates to a plant  
10 seed-specific expression vector containing the promoter and/or the intron and a transgenic plant transformed with the seed-specific expression vector.

Background Art

15 Fatty acids of plants are important components that make cell membranes and seed oils. In particular, microsomal oleic acid desaturase, which is present in the endoplasmic reticulum of plant cells, is an enzyme that catalyzes the conversion of monounsaturated fatty acid, oleic acid, at the sn-1 and sn-2 positions of phosphatidylcholine,  
20 into diunsaturated fatty acid, linoleic acid.

There have been reports about presence of genomic genes (*FAD2*) encoding the microsomal oleic acid desaturase in *Arabidopsis*, *Petunia*, and cotton. According to the reports, while one *FAD2* gene is present on the genome of *Arabidopsis*, two or more *FAD2* genes are  
25 present on each genome of *Petunia* and cotton [Okuley et al., 1994, Plant Cell; Verwoert et al., 2000, Biochemistry Society Transactions; Pirtie et al., 2001, Biochimica et Biophysica Acta]. In particular, it has been known that when two or more *FAD2* genes are present on a plant genome, at least one is involved in production of linoleic acid that  
30 accumulates in a seed oil. Until now, while the presence of the *FAD2* gene in the above-mentioned three plants has been reported, no studies on the functional analysis of its promoter have been reported.

WO 2004/044205

PCT/KR2003/002415

Recently, many attempts have been made to improve genetic characteristics of plants using genetic engineering technology. In this respect, expression of a useful foreign gene in a transgenic plant requires a promoter for the gene expression. For this, conventionally, a promoter derived from a cauliflower mosaic virus (CaMV35S) gene that induces gene expression at all tissues of a plant has been widely used. However, this promoter cannot induce seed-specific gene expression. Meanwhile, there have been reports about promoters that induce tissue-specific expression; for example, seed-specific expression [Plant Cell Technology, 1991, 3: 568-576], leaf- or flower-specific expression [Science, 1990, 250: 931-936], and root-specific expression [Plant Cell Technology, 1991, 3: 577-587]. However, the above seed-specific expression promoter induces gene expression during whole stages for seed development, not during a specific development stage.

In view of these problems, based on the fact that the microsomal oleic acid desaturase derived from sesame is specifically expressed during a seed development stage, the present inventors cloned the genomic gene encoding microsomal oleic acid desaturase and its promoter, inserted the promoter into a binary vector and a transient expression vector, and then introduced the vectors into the seeds of model plants, *Arabidopsis* and sesame. As a result, the present inventors found that the sesame microsomal oleic acid desaturase (*Si-FAD2*) promoter is a novel promoter that induces the expression of a foreign gene in a seed development stage-specific manner, and completed the invention.

Furthermore, a 127 bp active fragment of the promoter essential for the seed-specific expression has been identified. Also, the present inventors found that when the promoter is used together with an intron present on the *Si-FAD2* gene, the expression level of a foreign gene in a seed is enhanced by at least 40-fold, and completed the invention. In this regard, the intron is useful in high expression of a useful foreign gene.

WO 2004/044205

PCT/KR2003/002415

Therefore, the present invention provides a seed-specific, in particular, a seed development stage-specific expression promoter derived from the *Si-FAD2* gene and a 127 bp active fragment thereof.

The present invention also provides an intron that enhances the expression level of a foreign gene under the control of the above-described seed-specific expression promoter.

The present invention also provides a seed-specific expression vector containing the above-described promoter and/or intron and a transgenic plant transformed with the seed-specific expression vector.

The present invention also provides a method for expressing a foreign gene in a transgenic plant using the above promoter. This method is useful in producing a useful product in a seed development stage-specific manner or functionally modifying conventional product in seeds.

#### Disclosure of the Invention

Therefore, according to an aspect of the present invention, there is provided a seed-specific expression promoter comprising an active fragment of nucleotides -179 to -53 of SEQ ID NO: 3. Preferably, the present invention provides a seed-specific expression promoter comprising the nucleotide sequence as set forth in SEQ ID NO: 3.

The nucleotide sequence of SEQ ID NO: 3 comprises nucleotides that extend from position -660 to -1 from a transcription initiation site of the sesame microsomal oleic acid desaturase (*Si-FAD2*) gene which is expressed in a seed-specific manner. In this regard, it is expected that the nucleotide sequence contains a seed-specific transcription regulatory site as well as a TATA box and a CAAT-box which are common promoter sequences for recognition of the transcription initiation site of a plant gene. Also, nucleotides that extend from position -179 to -53 of SEQ ID NO: 3 correspond to the promoter active fragment (127 bp) essential for seed-specific expression. That is, when these nucleotides are deleted, promoter activity is greatly decreased.

WO 2004/044205

PCT/KR2003/002415

According to another aspect of the present invention, there is provided an intron for enhancement of gene expression, comprising nucleotides 149 to 1722 of SEQ ID NO: 1.

5 The nucleotides 149 to 1722 of SEQ ID NO: 1 corresponds to the intron present in 5'-untranslated region of the *Si-FAD2* gene. When the promoter is used together with the intron, the expression level of a foreign gene in a seed can be increased by at least 40-fold.

According to another aspect of the present invention, there is provided a seed-specific expression vector comprising the  
10 above-described promoter and/or intron.

Preferably, the seed-specific expression vector is constructed by inserting the promoter upstream of a foreign gene of a binary vector. The binary vector that can be used herein may be one that contains the left and right borders (BR and BL) of T-DNA capable of transforming a  
15 plant in the presence of the Ti-plasmid of *Agrobacterium tumefaciens*. For example, pGA series, pCG series, pCIT series, pGPTV series, pBECK2000 series, or pGreen series vectors may be used. It is preferable to use pBI series vectors (Clontech, America), easily commercially available in the pertinent art. The foreign gene that can  
20 be used herein may be a target gene or a reporter gene derived from an external source that is to be expressed in a plant of interest.

In the seed-specific expression vector of the present invention, the promoter is operably linked to a foreign gene contained in the binary vector. By "operably linked" is intended a functional linkage between a  
25 promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. For example, there is a pBin*Si-FAD2*-GUS expression vector (FIG. 4B) constructed by inserting the promoter into a pBI101 binary vector containing  $\beta$ -glucuronidase (GUS) gene which is a  
30 reporter gene. It is understood that the GUS reporter gene can be replaced with a desired foreign gene.

Furthermore, in the seed-specific expression vector of the present

WO 2004/044205

PCT/KR2003/002415

invention, the promoter and the intron are operably linked to a foreign gene contained in the binary vector. For example, there is a pSiW6-P2.4 expression vector (FIG. 9A) constructed by inserting the promoter and the intron into the pBI101 binary vector containing the GUS reporter gene. It is understood that the GUS reporter gene can be replaced with a desired foreign gene.

Preferably, the seed-specific expression vector of the present invention is constructed by inserting the promoter upstream of a foreign gene of a transient expression vector. Any transient expression vector can be used provided that it allows a foreign gene to be transiently expressed in plant tissues. For example, pBI221 [Mitsuhara et al., 1996], pMG221 [Maas et al., 1991], pUbiGUS [Christensen and Quail, 1996], or ACT1-D [McElroy et al., 1990] may be used. It is preferable to use pBI221 (Clontech, America) easily commercially available in the pertinent art. The foreign gene that can be used herein may be a target gene or a reporter gene derived from an external source that is to be expressed in a plant of interest.

In the seed-specific expression vector of the present invention, the promoter is operably linked to a foreign gene contained in the transient expression vector. For example, there is a pSiFAD2-GUS expression vector (FIG. 8A) constructed by inserting the promoter into a pBI221 transient expression vector containing the GUS reporter gene. It is understood that the GUS reporter gene can be replaced with a desired foreign gene.

According to another aspect of the present invention, there is provided a transgenic plant cell or plant transformed with the seed-specific expression vector.

When the seed-specific expression vector is a binary vector, a plant is transformed by floral dip method [Clough and Bent, 1998, The Plant Journal]. When the seed-specific expression vector is a transient expression vector, a plant is transformed by particle bombardment method [Lacorte et al., 1997, Plant Cell Reports]. The seed-specific

WO 2004/044205

PCT/KR2003/002415

expression vector can be used for transformation of all plants including dicot plant, monocot plant, sexual reproductive plant, and asexual reproductive plant. *Arabidopsis* and sesame (*Sesamum indicum*) are used herein.

5 According to yet another aspect of the present invention, there is provided a method for expressing a foreign gene in a transgenic plant transformed with the seed-specific expression vector.

The foreign gene may be any one that is wanted to be expressed in a plant. The foreign gene is positioned downstream of the promoter  
10 and/or the intron in the seed-specific expression vector. When needed, the expression of the foreign gene can be coupled with the expression of a reporter gene.

The present invention provides a seed-specific expression promoter of the *Si-FAD2* gene and a method for expressing a foreign  
15 gene in a transgenic plant using the promoter. Therefore, the present invention is useful in producing a useful product in a seed development stage-specific manner or functionally modifying a common product in a seed. Also, according to the present invention, the promoter can be used together with an intron for expression enhancement, thereby  
20 increasing the expression level of a foreign gene in a seed by at least 40-fold. Therefore, the present invention can be used in development of a transgenic plant which induces high expression of a foreign gene in a seed-specific manner.

25 Brief Description of the Drawings

FIG. 1 is the nucleotide sequence (SEQ ID NO: 1) of the gene encoding sesame microsomal oleic acid desaturase (*Si-FAD2*) and the amino acid sequence (SEQ ID NO: 2) deduced therefrom;

FIG. 2 is a diagram of *Si-FAD2* gene;

30 FIG. 3 is the nucleotide sequence (SEQ ID NO: 3) of the promoter of *Si-FAD2* gene;

FIG. 4A is a genetic map of a pBI101 binary vector containing

WO 2004/044205

PCT/KR2003/002415

$\beta$ -glucuronidase (GUS) gene used in a preferred embodiment of the present invention;

FIG. 4B is a diagram of a pBinSiFAD2-GUS expression vector constructed by inserting Si-FAD2 promoter into pBI101 binary vector  
5 containing GUS gene;

FIG. 4C is a polymerase chain reaction (PCR) result with vector gene-specific primers in *Agrobacteria* transformed with a pBinSiFAD2-GUS expression vector containing Si-FAD2 promoter or a pBI121 binary vector containing CaMV35S promoter;

10 FIG. 5 is a photograph of *Arabidopsis thaliana* transformed with *Agrobacteria* containing pBinSiFAD2-GUS or pBI121 vector;

FIG. 6 is a histochemical staining result for GUS activity in various tissues of *Arabidopsis thaliana* transformed with *Agrobacteria* containing pBinSiFAD2-GUS vector;

15 FIG. 7A is a graph showing relative GUS activity (%) in various tissues of *Arabidopsis thaliana* transformed with *Agrobacteria* containing pBinSiFAD2-GUS vector;

FIG. 7B is a graph showing relative GUS activity (%) in developing seeds of *Arabidopsis thaliana* transformed with *Agrobacteria* containing  
20 pBinSiFAD2-GUS or pBI121 vector;

FIG. 8A is a diagram of a pSiFAD2-GUS expression vector constructed by inserting Si-FAD2 promoter into a pBI221 transient expression vector containing GUS gene;

FIG. 8B is a histochemical staining result for GUS activity in  
25 developing seeds of sesame to which a pSiFAD2-GUS vector is introduced by particle bombardment.

FIG. 9A is a diagram of a pSiW6-P2.4 expression vector constructed by inserting the promoter and intron of Si-FAD2 gene into a pBI101 binary vector containing GUS gene, and pSiW6-F1, pSiW6-F2,  
30 pSiW6-F3, pSiW6-F4, and pSiW6-F5 expression vectors constructed by inserting whole or part of the promoter of the Si-FAD2 gene into the pBI101 binary vector containing the GUS gene;



WO 2004/044205

PCT/KR2003/002415

FIG. 9B is an electrophoretic photograph of the six expression vectors, pSiW6-P2.4, pSiW6-F1, pSiW6-F2, pSiW6-F3, pSiW6-F4, and pSiW6-F5 of FIG. 9A digested with *Hind*III and *Bam*HI restriction enzymes;

5        FIG. 10 is a photograph of *Arabidopsis thaliana* transformed with *Agrobacteria* containing the six expression vectors, pSiW6-P2.4, pSiW6-F1, pSiW6-F2, pSiW6-F3, pSiW6-F4, and pSiW6-F5 of FIG. 9A and a pBI121 binary vector;

10        FIG. 11 is a histochemical staining result for GUS activity in developing seeds, shells covering the seeds, and seedlings germinated from the seeds of the seven types of the *Arabidopsis thaliana* of FIG. 10;

FIG. 12A is a graph showing quantitative GUS activity (nmol/hr/mg protein) in developing seeds and shells covering the seeds of the seven types of the *Arabidopsis thaliana* of FIG. 10;

15        FIG. 12B is a graph showing quantitative GUS activity (nmol/hr/mg protein) in cotyledons of seedlings at 7 days after germination of the seeds of transformants selected after culturing the seeds of the seven types of the *Arabidopsis thaliana* of FIG. 10 in kanamycin (30  $\mu$ g/ml)-containing media; and

20        FIG. 13 is the nucleotide sequence (nucleotides -179 to -53 of SEQ ID NO: 3) of an active fragment of the seed-specific expression promoter of *Si*-FAD2 gene.

#### Best mode for carrying out the Invention

25        Hereinafter, the present invention will be described more specifically by Examples. However, the following Examples are provided only for illustrations and thus the present invention is not limited to or by them.

#### 30        Example 1: Cloning and sequencing of genomic gene encoding sesame microsomal oleic acid desaturase (*Si*-FAD2)

In order to obtain genomic clones of seed-specific *Si*-FAD2 gene,

WO 2004/044205

PCT/KR2003/002415

genomic DNA was extracted from young leaves of *Sesamum indium* [Dellaporta et al., 1984, in Molecular Biology of Plants]. The genomic clones of the *Si-FAD2* gene were isolated by polymerase chain reaction (PCR) using a pair of primers (forward primer W6-1: 5'-GACAAAATGGGAGCCGGAGGACGCATGT-3' and reverse primer W6-R1: 5'-CGGCTTCAGAACTTGTCTTGTACCAGA-3') designed from cDNA (GenBank Accession No: AF192486) disclosed by Jin et al. [2001, Plant Science]. The genomic clones thus isolated were cloned into a pGEM-T vector (Promega, America) followed by sequencing using ABI Bigdye cycle sequencing kit (PE Applied Biosystems, America). In comparison between the genomic clones and the cDNA, it was demonstrated that the genomic clones and the cDNA were identical except 9 bases (CGGCACGAG) from the 5'-end of the cDNA. The genomic clones of the *Si-FAD2* gene contained a single bulky intron containing 1,574 bp in the 5'-untranslated region (UTR). However, there was no intron in the coding region. The nucleotide sequence (SEQ ID NO: 1) of the *Si-FAD2* gene and the amino acid (SEQ ID NO: 2) deduced therefrom are shown in FIG. 1. The first base "A" is a translation initiation base, and bases in italic type (*GT-AG*) are those of the intron contained in the 5'-UTR.

#### Example 2: Sequencing of 5'-upstream region of *Si-FAD2* gene

In order to clone the promoter of the *Si-FAD2* gene, inverse PCR was performed using the genomic DNA of the *Si-FAD2* gene digested with *NdeI* restriction site. Generally, inverse PCR has been frequently used in detection of unknown regions such as a promoter [Digeon et al., 1999, Plant Molecular Biology]. For inverse PCR, the genomic DNA was digested with *NdeI* restriction enzyme and self-ligated by T4 DNA ligase (BM, Germany) to prepare a template DNA. Inverse PCR was performed in two steps to obtain specific PCR products. First, the first PCR was performed using gene-specific primers (W6-1: 5'-GACAAAATGGGAGCCGGAGGACGCATGT-3' and W6-R6:

WO 2004/044205

PCT/KR2003/002415

5'-GGGGGCACGTTACCTGAAAAGTTGGAAG-3') to obtain a first PCR product. Then, the second PCR was performed using the first PCR product as a template with inner primers (W6-2: 5'-GGCTTTGGGACGAAGACTTCGTCACGCT-3' and W6-R7: 5'-CGCGTGAAAGCACTTCTGCGGAAGCGC-3'). As a result, the 5'-upstream region (about 750 bp, -660 to +90) of the *Si-FAD2* gene was obtained. FIG. 2A shows a diagram of the *Si-FAD2* gene. Here, SiW6F1, SiW6R1, W6R3, W6R5, and W6R1 indicate the positions of primers as used in FIG. 2B.

10 In order to determine whether the obtained fragment is the 5'-upstream region of the *Si-FAD2* gene, PCR was performed using the gene-specific primers (SiW6F1, SiW6R1, W6R1, W6R3, and W6R5) designed from the 5'-end of the 750 bp fragment and the internal region of the *Si-FAD2* gene. As a result, it was demonstrated that the 750 bp  
15 fragment is the 5'-upstream region of the *Si-FAD2* gene. FIG. 2B shows a PCR result of the genomic DNA of sesame leaf *FAD2* gene using the primers of FIG. 2A. Here, M1 and M2 represent DNA size markers. Lane 1 is a PCR product obtained using the SiW6F1 (5'-CCGAAGCTTCATATGTGAAATGTAATGGAAAATGCGAC-3') and  
20 the SiW6R1 (5'-CTTGGATCCTTGAAGGAGAAATCGCGTGAAAGCAC-3'). Lane 2 is a PCR product obtained using the SiW6F1 and the W6R1 (5'-CGGCTTCAGAACTTGTTCTTGTACCAGA-3'). Lane 3 is a PCR product obtained using the SiW6F1 and the W6R5 (5'-GGAGAACGAACGGCTGACGGATCTCTCG-3'). Lane 4 is a PCR  
25 product obtained using the SiW6F1 and the W6R3 (5'-GGCTTTGGGACGAAGACTTCGTCACGCT-3'). The 5'-upstream region of the *Si-FAD2* gene was sequenced. The nucleotide sequence (SEQ ID NO: 3) of the promoter region of 1 to 660 bp long upstream  
30 from a transcription initiation site is shown in FIG. 3.

### Example 3: Identification of transcription initiation site of *Si-FAD2*

WO 2004/044205

PCT/KR2003/002415

gene

In order to determine the activity of the seed-specific promoter of the *Si-FAD2* gene, the transcription initiation site of the gene must be identified. For this, cRACE (circular first-strand cDNA-mediated rapid amplification of cDNA ends) was performed according to Maruyama et al  
5 method [Nucleic Acid Research, 1995]. First, total RNA was isolated from sesame seeds. Then, cDNA was synthesized with a gene-specific primer (5'-GGTAGCAGTATGGGGATGGCAGCAGATGGAAGTA) which was phosphorylated by T4 polynucleotide kinase (Takara) and reverse  
10 transcriptase (BM), and the 5'- and 3'-ends of the cDNA were ligated with T4 RNA ligase (NEB). PCR was performed using the resultant cDNA as a template with gene-specific primers (W6-5: 5'-GAAGAACCCCTCCAACGGGTGCC-3' and W6-R9: 5'-CCGATCACATCGCAAGTGCGATACACCTG-3'). As a result, it was  
15 demonstrated that the first base "A" of FIG. 1 is a transcription initiation base.

Example 4: Analysis of *Si-FAD2* promoter activity in *Arabidopsis*

In order to analyze the *Si-FAD2* promoter activity in *Arabidopsis*, a  
20 pBin*Si-FAD2*-GUS expression vector containing the cloned *Si-FAD2* promoter and GUS reporter gene was constructed. First, PCR was performed using sesame genomic DNA as a template with a pair of primers, the *SiW6F1* (5'-CCGAAGCTTCATATGTGAAATGTAATGGAAAATGCGAC-3') and  
25 the *SiW6R1* (5'-CTTGGATCCTTGAAGGAGAAATCGCGTGAAAGCAC-3'). As a result, the promoter region (-660 to +141) of the *Si-FAD2* gene was obtained. The promoter region and a pBI101 binary vector (Cat. # 6017-1, Clontech, America) of which genetic map is shown in FIG. 4A  
30 were digested with the same restriction enzymes *HindIII* and *BamHI*. The digested promoter region and pBI101 binary vector were ligated using T4 DNA ligase to construct the pBin*Si-FAD2*-GUS expression

WO 2004/044205

PCT/KR2003/002415

vector. FIG. 4B shows a diagram of the pBinSiFAD2-GUS expression vector in which the promoter region was cloned into the *HindIII*-*Bam*HI restriction sites of the pBI101 binary vector. Here, GUS is a reporter gene encoding  $\beta$ -glucuronidase (which can be replaced with a foreign gene to be expressed). NPTII is a kanamycin resistance marker gene encoding neomycin phosphotransferase II. Nos-pro and Nos-ter are respectively the promoter and terminator for plant expression of the NPTII.

The pBinSiFAD2-GUS expression vector thus constructed was introduced into *Agrobacterium tumefaciens* C58C1 [Suh et al., 2002, Molecules and Cells] by freeze-thaw method [An, G. 1987, Methods in Enzymology]. According to the freeze-thaw method, the *Agrobacteria* were suspension-cultured in YEP media until O.D = 0.5 followed by resuspending in 20 mM  $\text{CaCl}_2$  solution. The suspension thus obtained was mixed with the pBinSiFAD2-GUS expression vector and incubated in liquid nitrogen for 1 minutes and then at 37°C for 2 minutes. FIG. 4C shows a PCR result of the *Agrobacteria* transformed with the pBinSiFAD2-GUS expression vector of FIG. 4B and pBI121 binary vector (Cat. # 6018-1, Clontech, America) containing CaMV35S promoter using vector gene-specific primers. Here, M is a DNA size marker, N is a PCR product of pBI121 binary vector-containing *Agrobacteria* using the SiW6F1 and SiW6R1 (negative control), P is a PCR product of pBinSiFAD2-GUS expression vector-containing *Escherichia. coli* using the SiW6F1 and SiW6R1 (positive control); SS is a PCR product of pBinSiFAD2-GUS expression vector-containing *Agrobacteria* using the SiW6F1 and SiW6R1; SN is a PCR product of pBinSiFAD2-GUS expression vector-containing *Agrobacteria* using a pair of primers (forward primer: 5'-GAGGCTATTCGGCTATGACTG-3' and reverse primer: 5'-ATGGGAGCGGCGATACCGTA-3') in npt II gene (kanamycin resistance marker gene), and V is a PCR product of pBI121 binary vector-containing *Agrobacteria* using the primers (forward primer: 5'-GAGGCTATTCGGCTATGACTG-3' and reverse primer:

WO 2004/044205

PCT/KR2003/002415

5'-ATGGGAGCGGCGATACCGTA-3')) in the npt II gene. As a result, when the nptII primers were used, PCR products of about 700 bp long were obtained. On the other hand, when the gene-specific primers, the SiW6F1 and the SiW6R1 were used, PCR products of about 750 bp long were obtained. The result of FIG. 4C demonstrates that the pBinSiFAD2-GUS and pBI121 vectors were introduced into the *Agrobacteria*.

After the *Agrobacteria* containing the pBinSiFAD2-GUS and pBI121 vectors were shaking-cultured at 28°C for 2 days, the cultures were inoculated onto the pistils of *Arabidopsis thaliana* (ecotype Columbia) before flower opening by floral dip method [Clough and Bent, 1998, The Plant Journal]. FIG. 5 shows the *Arabidopsis thaliana* transformed with the *Agrobacteria* containing the pBinSiFAD2-GUS and pBI121 vectors of FIG. 4C. Here, a is the *Arabidopsis thaliana* transformed with the pBI121 vector-containing *Agrobacteria* and b is the *Arabidopsis thaliana* transformed with the pBinSiFAD2-GUS vector-containing *Agrobacteria*.

Transgenic plants were selected on the MS media containing kanamycin (30 µg/ml). The GUS activity under the control of the Si-FAD2 promoter in various tissues of the transgenic plants was analyzed by a histochemical staining method and an enzymatic method. FIG. 6 is a histochemical staining result for GUS activity in various tissues of *Arabidopsis thaliana* transformed with the *Agrobacteria* containing the pBinSiFAD2-GUS vector under the control of the Si-FAD2 promoter. Here, (A) is *Arabidopsis thaliana* untreated, (B) is *Arabidopsis thaliana* transformed with pBI121 vector-containing *Agrobacteria*, and (C) is *Arabidopsis thaliana* transformed with pBinSiFAD2-GUS vector-containing *Agrobacteria*.

FIG. 7A is a graph showing relative GUS activity (%) in various tissues of *Arabidopsis thaliana* transformed with *Agrobacteria* containing the GUS gene-containing binary vectors under the control of the Si-FAD2 promoter. Here, the minimal GUS activity was defined as 1%. Seed

WO 2004/044205

PCT/KR2003/002415

specimens were extracted depending on development stages (seeds 1: 5 to 7 days after flowering, seeds 2: 10 to 15 days after flowering, and seeds 3: 20 to 25 days after flowering). FIG. 7B is a graph showing relative GUS activity (%) in developing seeds of *Arabidopsis thaliana* transformed with *Agrobacteria* containing the pBinSiFAD2-GUS or pBI121 binary vector. Here, the minimal GUS activity was defined as 1%. As seen from FIGS. 6, 7A, and 7B, the GUS gene was expressed in an *Arabidopsis thaliana* seed-specific manner under the control of the Si-FAD2 promoter. In particular, the seeds that had developed for 10 to 25 days after flowering exhibited the highest level of GUS activity. These results demonstrate that the Si-FAD2 promoter can induce the expression of a foreign gene in a seed-specific manner, in particular a seed development stage-specific manner.

Example 5: Analysis of Si-FAD2 promoter activity in sesame seeds

In order to analyze the activity of the Si-FAD2 promoter in sesame seeds, a pSiFAD2-GUS transient expression vector containing the cloned Si-FAD2 promoter and the GUS reporter gene was constructed. First, the Si-FAD2 promoter region (-660 to +141) obtained in Example 4 was digested with *Hind*III and *Bam*HI restriction enzymes. On the other hand, the CaMV35S promoter was removed from pBI221 (Cat. # 6019-1, Clontech, America) transient expression vector by using *Hind*III and *Bam*HI restriction enzymes. The Si-FAD2 promoter was inserted into the CaMV35S promoter site of the pBI221 transient expression vector by using T4 DNA ligase. As a result, the pSiFAD2-GUS transient expression vector containing the Si-FAD2 promoter instead of the CaMV35S promoter was constructed. FIG. 8A shows a diagram of the pSiFAD2-GUS transient expression vector. Here, GUS is a reporter gene encoding  $\beta$ -Glucuronidase (which can be replaced with a foreign gene to be expressed). Nos-ter is the terminator for expression of nopaline synthase gene.

WO 2004/044205

PCT/KR2003/002415

Plasmid DNA from the p*Si*/FAD2-GUS transient expression vector was purified using plasmid midi kit (Qiagen, America) and coated with 1.6  $\mu$ m gold particles (Bio-rad, America). 2 ug of the plasmid DNA was mixed with 500 ug of the gold particles for one shot of particle bombardment (PDS1000/He gun, Bio-rad). The plasmid DNA mixed with the gold particles was bombarded under 1,100 psi (He gas) and incubated at 27°C for 18 hours. The activity of the GUS, a foreign gene, under the control of the *Si*-FAD2 promoter, was analyzed by a histochemical staining method. FIG. 8B is a histochemical staining result for GUS activity in sesame developing seeds to which the p*Si*/FAD2-GUS vector of FIG. 8A is introduced by particle bombardment. Here, **a** is the pBI221 vector containing no promoters, **b** is the pBI221 vector containing the CaMV35S promoter, and **c** is the p*Si*/FAD2-GUS vector containing the *Si*-FAD2 promoter. As seen from FIG. 8B, the *Si*-FAD2 promoter can induce the expression of a foreign gene such as GUS in sesame developing seeds.

Example 6: Construction of binary vectors containing the active fragment of *Si*-FAD2 promoter and expression enhancing intron of *Si*-FAD2 gene

In this Example, the active fragment of the *Si*-FAD2 promoter involved in seed-specific expression was identified and the effect of the intron present on the *Si*-FAD2 gene on expression of a foreign gene in a transgenic plant was analyzed. For this, first, gene-specific primers containing *Hind*III (AAGCTT) and *Bam*HI (GGATCC) restriction sites as presented in Table 1 were designed using the nucleotide sequence of the *Si*-FAD2 promoter.

Table 1

30

Primer type	Name	Nucleotide sequence	Tm
Forward	SiW6F1	5'-CCGAAGCTTCATATGTGAAATGTAATGAAAATGCGAC-3'	72°C



WO 2004/044205

PCT/KR2003/002415

primer	SiW6F2	5'-CCGAAGCTTGGGACATGCCACATTATGTGG-3'	67°C
	SiW6F3	5'-CCGAAGCTTGTCTTAACCAGGTTTGAACAACC-3'	67°C
	SiW6F4	5'-CCGAAGCTTGGAAATGTGCACACTCCATGTG-3'	67°C
	SiW6F5	5'-CCGAAGCTTGGGCCCCCTCTCAGACAGG-3'	71°C
Reverse primer	SiW6R1	5'-CTTGGATCCTTGAAGGAGAAATCGCGTGAAAGCAC-3'	75°C
	SiW6R3	5'-CAAGGATCCGTCAAGCCGCCCAATTTAC-3'	68°C

Next, PCR was performed by using sesame genomic DNA as a template with the above primers to obtain PCR products containing an original promoter region and a promoter region with partial deletion from the 5'-end. The PCR products thus obtained were digested with *Hind*III and *Bam*HI and then inserted into the *Hind*III-*Bam*HI restriction sites of pBI101 binary vectors (FIA. 4A, Cat. # 6017-1, Clontech, America) using T4 DNA ligase to construct binary vectors, pSiW6-P2.4, pSiW6-F1, pSiW6-F2, pSiW6-F3, pSiW6-F4, and pSiW6-F5 (FIG. 9A). FIG. 9A shows a diagram of the pSiW6-P2.4 binary vector containing the promoter and intron of the *Si*-FAD2 gene. Here, GUS is a reporter gene encoding  $\beta$ -glucuronidase (which can be replaced with a foreign gene to be expressed). NPTII is a kanamycin resistance marker gene encoding neomycin phosphotransferase II. Nos-pro and Nos-ter are respectively the promoter and terminator for plant expression of the NPTII. The GUS reporter gene is expressed in a plant under the control of the inserted promoter and the Nos terminator (Nos-ter).

The binary vectors thus constructed were introduced into *Escherichia. coli* and plasmid DNA was extracted by an alkaline lysis method [Sambrook et al., 2001]. The extracted plasmid DNA was digested with *Hind*III and *Bam*HI followed by 0.7% agarose gel electrophoresis (FIG. 9B). FIG. 9B shows an electrophoretic result of the six binary vectors digested with *Hind*III and *Bam*HI. Here, pSiW6-P2.4 is a binary vector containing the promoter and intron of a PCR product amplified with the gene-specific primers, SiW6F1 (-660 bp) and SiW6R3 (+1743 bp). pSiW6-F1 is a binary vector (identical to pBinSiFAD2-GUS binary vector) containing the promoter of a PCR

WO 2004/044205

PCT/KR2003/002415

product amplified with the gene-specific primers, SiW6F1 (-660 bp) and SiW6R1 (+141 bp). pSiW6-F2 is a binary vector containing the promoter of a PCR product amplified with the gene-specific primers, SiW6F2 (-547 bp) and SiW6R1 (+141 bp). pSiW6-F3 is a binary vector  
5 containing the promoter of a PCR product amplified with the gene-specific primers, SiW6F3 (-346 bp) and SiW6R1 (+141 bp). pSiW6-F4 is a binary vector containing the promoter of a PCR product amplified with the gene-specific primers, SiW6F4 (-179 bp) and SiW6R1 (+141 bp). pSiW6-F5 is a binary vector containing the promoter of a  
10 PCR product amplified with the gene-specific primers, SiW6F5 (-52 bp) and SiW6R1 (+141 bp).

#### Example 7: Transformation of *Arabidopsis* with binary vectors

The six binary vectors (pSiW6-P2.4, pSiW6-F1, pSiW6-F2, pSiW6-F3, pSiW6-F4, and pSiW6-F5) constructed in Example 6 and the  
15 pBI121 binary vector (Cat.# 6018-1, Clontech, America) were introduced into *Agrobacterium tumefaciens* C58C1 [Suh et al., 2002, Molecules and Cells] by freeze-thaw method [An, G. 1987, Methods in Enzymology]. According to the freeze-thaw method, the *Agrobacteria* were  
20 suspension-cultured in YEP media until O.D = 0.5 followed by resuspending in 20 mM CaCl<sub>2</sub> solution. The suspension thus obtained was mixed with each of the seven binary vectors and incubated in liquid nitrogen for 1 minutes and then at 37°C for 2 minutes. The transformed  
25 *Agrobacteria* were shaking-cultured at 28°C for 2 days, and the cultures were inoculated onto the pistils of *Arabidopsis thaliana* (ecotype Columbia) before flower opening by floral dip method [Clough and Bent, 1998, The Plant Journal].

FIG. 10 is a photograph of *Arabidopsis thaliana* transformed with the *Agrobacteria* containing the six binary vectors of FIG. 9B and the  
30 pBI121 binary vector.

#### Example 8: Histochemical staining and enzymatic analysis for

WO 2004/044205

PCT/KR2003/002415

GUS activity in transformed *Arabidopsis thaliana*

Seeds were harvested from the seven independent *Arabidopsis* transgenic lines constructed in Example 7 and spread on MS media containing kanamycin (30  $\mu\text{g/ml}$ ) to select kanamycin resistance transformants. GUS activity in the individual tissues of the selected transformants was analyzed by a histochemical staining method and an enzymatic method. In order to stain the individual tissues of the transformants, the individual tissues were immersed in a solution containing 1 mM X-glu (5-bromo-4-chloro-3-indolyl- $\beta$ -glucuronide), 100 mM sodium phosphate (pH 7.0), 10 mM EDTA, 0.5 mM potassium ferricyanide, 0.5 mM potassium ferrocyanide, and 0.1% Triton X-100) and incubated at 37°C for 12 hours. After removal of the solution, the stained tissues were rinsed with 100% ethanol until chlorophyll was exuded from the tissues. For the quantitative analysis of the GUS activity, the individual tissues were homogenized in a solution containing 50 mM sodium phosphate (pH 7.0), 10 mM EDTA, 0.1% Triton X-100, 0.1% sodium lauroylsarcosine, and 10 mM  $\beta$ -mercaptoethanol according to Jerrerson et al method [EMBO J. 6: 3901-3907, 1987] followed by centrifugation (12,000 g) to obtain supernatants. The supernatants were mixed with 1 mM MUG (4-methylumbelliferyl glucuronide) and incubated at 37°C. The reaction was terminated by adding 0.2 M  $\text{Na}_2\text{CO}_3$ . Fluorescence of the resultant reactions was measured at 365 and 455 nm by a fluorometer and plotted on the standard curve generated using the MUG standard to quantify the GUS activity.

FIG. 11 is a histochemical staining result for GUS activity in developing seeds (seeds 1: 5 to 7 days after flower opening, seeds 2: 10 to 15 days after flower opening, seeds 3: 20 to 25 days after flower opening), shells covering the seeds, and seedlings germinated from the seeds of the *Arabidopsis thaliana* transformed with *Agrobacteria* containing the seven binary vectors. CaMV35S is *Arabidopsis thaliana*

WO 2004/044205

PCT/KR2003/002415

transformed with pBI121 binary vector. SiW6P2.4, SiW6F1, SiW6F2, SiW6F3, SiW6F4, and SiW6F5 are *Arabidopsis thaliana* transformed with pSiW6-P2.4, pSiW6-F1, pSiW6-F2, pSiW6-F3, pSiW6-F4, and pSiW6-F5, respectively. According to the results of FIG. 11, the  
5 Si-FAD2 promoter with a deletion of about 150-200 bp long from the 5'-end region also induced the expression of a foreign gene in a seed-specific manner. In addition, when the intron in the 5' UTR of the Si-FAD2 gene was inserted downstream of the Si-FAD2 promoter region (-660/+141) (SiW6P2.4, -660/+1743), GUS expression was detected in  
10 the shells and the cotyledons of the seedlings, in addition to the developing seeds.

FIG. 12A shows GUS activity in developing seeds (seeds 1: 5 to 7 days after flowering, seeds 2: 10 to 15 days after flowering, seeds 3: 20 to 25 days after flowering) and shells covering the seeds of the seven  
15 independent *Arabidopsis* transgenic lines obtained in FIG. 11. The GUS activity was analyzed in 10 transformants (T1 plant) per each of the seven transgenic plant constructs. As shown in FIG. 12A, the GUS activity of the transgenic plant constructs containing the Si-FAD2 promoter with a gradual deletion of 150-200 bp long from the 5'-end  
20 region was analyzed. As a result, SiW6F1 (-660/+141), SiW6F2 (-547/+141), SiW6F3 (-346/+141), and SiW6F4 (-179/+141) constructs exhibited the GUS activity of more than about 4 to 10-fold in the seeds 2, as compared to SiW6F5 construct (-52/+141). It can be seen from the result that the nucleotide sequence of -179 to -53 long in the Si-FAD2  
25 promoter is an essential site that induces seed-specific expression. Also, SiW6P2.4 construct (-660/+1743) containing the intron exhibited the GUS activity of more than about 40-fold in the seeds 3 and more than 10-fold in the shells, as compared to the SiW6F1 construct (-660/+141) containing no intron. From the result, it can be seen that  
30 the Si-FAD2 promoter in the presence of the intron of the 5' UTR can greatly increase the expression level of a foreign gene in developing seeds and shells.

WO 2004/044205

PCT/KR2003/002415

FIG. 12B shows GUS activity in seedlings (T2 plant) at 7 days after germination of the seeds of kanamycin (30  $\mu$ g/ml) resistant transformants selected from the *Arabidopsis* transformants of FIG. 11. The GUS activity was analyzed in 5 transformants (T2 generation) per each of the seven transgenic plant constructs. GUS activity was not significantly observed in the seedlings of the SiW6F1, SiW6F2, SiW6F3, SiW6F4, and SiW6F5 constructs containing only the *Si-FAD2* promoter. On the other hand, the SiW6P2.4 construct containing the promoter and intron of the *Si-FAD2* gene exhibited about more than 30-fold increase of GUS activity in the cotyledons thereof, as compared to the SiW6F1 construct containing no intron. From these results, it can be seen that while the *Si-FAD2* promoter induces the expression of a foreign gene in a developing seed, the intron of the *Si-FAD2* gene can enhance the expression of a foreign gene in a seed-, shell-, and seedling cotyledon.

15

#### Industrial Applicability

As is apparent from the above descriptions, the present invention provides the nucleotide sequence of sesame microsomal oleic acid desaturase (*Si-FAD2*) gene and the *Si-FAD2* promoter. The *Si-FAD2* promoter induces the expression of a foreign gene in a plant in a seed development stage-specific manner. The present invention also provides a method for expressing a foreign gene in a transgenic plant using the *Si-FAD2* promoter. Therefore, the present invention is useful in producing a useful product in a seed-specific manner or functionally modifying a common product in a seed. Also, according to the present invention, the promoter can be used together with the intron for expression enhancement, thereby increasing the expression level of an inserted gene in a seed by at least 40-fold. Therefore, the present invention can be used in development of a transgenic plant which induces large-scale expression of a foreign gene in a seed-specific manner.

30

WO 2004/044205

PCT/KR2003/002415

What is claimed is:

1. A seed-specific expression promoter comprising an active fragment of nucleotides -179 to -53 of SEQ ID NO: 3.
- 5 2. The seed-specific expression promoter of claim 1, which comprises the nucleotide sequence as set forth in SEQ ID NO: 3.
3. An intron for enhancement of gene expression comprising nucleotides 149 to 1722 of SEQ ID NO: 1.
- 10 4. A seed-specific expression vector comprising the promoter of claim 1 and/or the intron of claim 3.
5. The seed-specific expression vector of claim 4, which is  
15 constructed by inserting the promoter upstream of a foreign gene of a binary vector.
6. The seed-specific expression vector of claim 5, which is a pBinSiFAD2-GUS expression vector as shown in FIG. 4B.
- 20 7. The seed-specific expression vector of claim 5, which is a pSiW6-P2.4 expression vector as shown in FIG. 9A.
8. The seed-specific expression vector of claim 4, which is  
25 constructed by inserting the promoter upstream of a foreign gene of a transient expression vector.
9. The seed-specific expression vector of claim 8, which is a pSiFAD2-GUS expression vector as shown in FIG. 8A.
- 30 10. A transgenic plant transformed with the seed-specific expression vector of any one of claims 4 through 9.

**WO 2004/044205****PCT/KR2003/002415**

11. A method for expressing a foreign gene in a transgenic plant transformed with the seed-specific expression vector of any one of claims 4 through 9.

5

FIG. 1a

[illegible]



10/534128

WO 2004/044205

PCT/KR2003/002415

2/12

FIG. 1b

ATGGGAGGCGGGAGGACGCGATGCTGTGATCCAAAGACGAAAGAGCGAACAAAGAAAGAACCCG  
M G A G G R M S D P T T K D E Q K K N P  
CTCCAAACGGGTGCGCTTACGCAAAAGCGCTCCATTGACACTCGGTGACATCAAGAAAGCGCAT  
L Q R U P Y A K P P F T L G D I K K A I  
CGACCACACTGCGITCGAGAGATCGGTGAGCGCGITCGITCTCTATGTCGTTTACGATCTC  
P P H C F E R S U S R S F S Y U U Y D L  
GTCAITGTTTTCTCTCTACTACATTCGCGATCTCTACTTCCATCTGCTGCGCATCCGCA  
U I U F L L Y Y I A T S Y F H L L P S P  
TACTGCTACCTAGCTTGGCGGATTTACTGGGCTGTACAAAGGCTGCGTTTGGCAGCGGAATC  
Y C Y L A H P I Y M A U Q G C U C T G I  
TGGGTCATTGCCCCATGAATGTEGGCCACCATGCGATTGAGCGATTACCGATGGCTTACGAC  
H U I A H E C G H H A F S D Y Q U L D D  
ACAGTGGCGCTCATCTGCGACTCTGCGCTGCTCGGTGCGGATTTTCTCATGGAATACAGC  
T U G L I L H S A L L U P Y F S H K Y S  
CAGCGCGCGCCACCACTCCAAACACTGGATCGCTTGGCGGTGACGAAGCTCTCTGTCCTCAAG  
H R R H H S N T G S L E R D E U F U P K  
CCAAAAITCAGAGTCTCGTGGTACTCCAAATACTTGAACAATCCACTTGGCAGAGTCATC  
P K S R U S M Y S K Y L N N P L G R U I  
ACACITGCTGGTTACTCTTACTCTCGGATGGCGCTTACTTGGCTGTTAATGCTCTGCGC  
T L U U T L T L G W P L Y L L F N U S G  
AGGCGTTACAAACGCTTTTGCATGCCAOTTTGACCCATATGGTCCAATATATATGACCGT  
R P Y N R F A C H F D P Y G P I Y N D R  
GAGAGACTTCAAACTCTCATCTCGATGCTGGTATATTTGCTGCTGATGCTGCTTTAT  
E R L Q I F I S D A G I I A A U D U L Y  
CGTGTGCTTTGGTCAAGGCTTGGCTTGGCTGCTATGCTTTATGGGCTAGCGTTACTC  
R U A L U K G L A W L U C U Y G U P L L  
ATTGTCAGCGCTTCTCTTTTGTATGATGATCTCTTCAAGCACATCAGCGTTCTGTTGCG  
I U N G F L U L I T F L Q H T H P S L P  
CACTATGATTTCTCGAGTGGGCTGGCTAAGGGGAGCTCTTGCACCTGTGACAGAGAT  
H Y D S S E W D W L R G A L A T U D R D  
TAGGGGTGCTAAATAGGTGCTTCCATACATCAGATACGACGCTGACTCACCAGCTT  
Y G U L N K U F H N I T D T H U T H H L  
TTCTCAAGGATGCCACATTACCATGCAATGGAGGCACTAAGGCAATCAAGGCCATACIG  
F S T M P H Y H A M E A T K A I K P I L  
GGCCAGTATATCAGTTTGTATGGAACCGCGTTTACAGGCGGATGTCGAGGAGGCAAG  
G Q Y Y Q F D G T P F Y K A M W R E A K  
GAATGCTGTATGTGAGCGAGACGAGAGTACTCCAGACAAGGCTGATTTCTGGTACAG  
E C L Y U E P D E S T P D K Q U F W Y K

10/534128

WO 2004/044205

PCT/KR2003/002415

3/12

FIG. 1c

AACCAAGTTCGAGGCCGGAATAACATGTTGGTTAGTGAAGATGGCGTCTTCTTATTTTGTCC

N K F -

TATCGAGATGGAGGAACATCATCATTTTTGTTTTCTTCTTATAAGATGCGTCCCTTGT

TAGTGTATTCCTGCACTTAATAAATAAAGCTCTACCCGAAGCCTGCTGCTGCTGCTG

GGATTCTAGTTCTGCAATAAATTGTCAGTTTAGTG

10/534128

WO 2004/044205

PCT/KR2003/002415

4/12

FIG. 2a



FIG. 2b




FIG. 3

CATATGTGAATGTAAATGGAAAAATGCGACAAGAATTGCAATAGAGAAAAATCCAATTTGCAGAGATT  
 CATGAAAAGAATTTGTACAATAGCATATATATGTTAAATGAAATGGGACATGCCACATTATGTGG  
 AATAAAAAAGACAATTTGCTTGGAAATTAATTATAGAAATAAATGTGTTACATTTAAATATGTGATTAAAT  
 CACTTTTTTTGAATTGTACATCTATCACATGACAAGTTCATTATATTTGACATATAATTTGTTTTATG  
 TCTAGTCAAGCCTAATTAAATTTCTCGGAAGACAAAAATTTTTTTGTCCTAACCAGGTTTGAACAA  
 CCAAA CAAATCACAAGCAGGTGTATCGCACTTGCATGTGATCGGTCACCTTTTCTAAATTTGTACA  
 TCATTACACGACAACGTATTTGTGCTCCAAGTTCAAATTGAGTGGGTTGGAGCTATAATTTCTTGT  
 AACACACAAATGTGGAAATGTGCACACTCCATGTGGGCCAATGAGCGGATGACACGTGGCGGGCACTT  
 ACCTCGTTACGTTGAGGCAATGATGAAGGGGGATCTCTTGAAGGTGGAGGGGTGGGGGGCGGGGGTTG  
 GGGGGGGGCCCCCTCCTCAGACAGGTCTATA TTATGAGACCTCGTAAGGCAGAACGC

10/534128

WO 2004/044205

PCT/KR2003/002415

5/12

FIG. 4a





FIG. 4b



10/534128

WO 2004/044205

PCT/KR2003/002415

6/12

FIG. 4c




FIG. 5



10/534128

WO 2004/044205

PCT/KR2003/002415

7/12

FIG. 6





FIG. 7a



10/534128

WO 2004/044205

PCT/KR2003/002415

8/12

FIG. 7b





FIG. 8a



WO 2004/044205

10/534128

PCT/KR2003/002415

9/12

FIG. 8b





FIG. 9a





10/534128

WO 2004/044205

PCT/KR2003/002415

10/12

FIG. 9b




FIG. 10



10/534128

WO 2004/044205

PCT/KR2003/002415

11/12

FIG. 11





FIG. 12a



10/534128

WO 2004/044205

PCT/KR2003/002415

12/12

FIG. 12b




FIG. 13

-179 GGAATGTGCACACTCCATGTGGGCCAATGAGCGGATGACACGTGGCGGG  
CAACTTACCTCGTTACGTTGAGGCATGCATGAAAGGGGGATCTCTTGAGGTGGA  
GGGGTGGCGCGGGGTTGGGGGG -53

JC06 Rec'd PCT/PTO 06 MAY 2005

WO 2004/044205

PCT/KR2003/002415

## SEQUENCE LISTING

<110> KOREA CHUNGANG EDUCATIONAL FOUNDATION

5 <120> Plant seed-specific expression promoter derived from sesame and seed-specific expression vector comprising the promoter

<130> PX208590V

10 <150> KR 10-2002-0069589  
<151> 2002-11-11

<160> 3

15 <170> KopatentIn 1.71

<210> 1  
<211> 3102  
<212> DNA  
20 <213> Sesamum indicum

<220>  
<221> 5'UTR  
<222> (1)..(1746)

25 <220>  
<221> intron  
<222> (149)..(1722)

30 <220>  
<221> 3'UTR  
<222> (2899)..(3102)

<400> 1

35 aagccactc aaatatttca ccaccaccac caccagaaca ttcagaaaca agaaataaac 60

acacacacac tataaaacag ttcttgcgaa agaaggaaag cgcttcgca gaagtgcctt 120

cacgcgattt ctcttccaa gtttcaggt aacgtgcccc cttttctctt ctctctatt 180

40 ctcttttctc ataattcatg atcaatcttt gagtattttg gtgtttgtgt gtctcaagaa 240

aaccgcattt ttattttctt gcaatggtgt ctttatttcc tgcgttttt ttcagctatt 300

45 aatgttcttt tgatgtagat gaggtttaat cgtatgttct tgagctgcat tacctgatga 360

ttcatggatc tgaggaatgt atcgcatitt ttatttttgt ttatttttt ggtgggcttt 420

ccaagaaga atctatttgg ggttattctt gtgtggtttg gtgcaaactt ttggatttta 480

50 cgcagtattg gtgtctggac cacatgattg tgcattttat atttgatttt tgcctttatc 540

WO 2004/044205

PCT/KR2003/002415

	tttgtatgca tgtgggatgc aggaagaaaa aactgtggta aatgtctttg aagagattga	600
	tttagcatat atacaagggt gcctgggctt cagttttgat gattttgatg tacattgtgg	660
5	agatttgatg ggttgcattg ggctcaaatc ttcttgaag atttgtttt tgtccaaaaa	720
	atttgggatt ttccacttt tattgaacag tagatctttt cctgtttcaa cccaaaagtt	780
10	atttcggttt gaagttttac atcatagata taatiagtaa taaatttcgg ttaggtccgt	840
	aaagaatcat taattacatc aattaatatt gttaaatga caaaaagagg gaatttatgg	900
	tgatatctat gaagccatgc tatgcctggc tggaattccg tcgatgaaaa agacagattc	960
15	cgggtgtgtg tagatttcac tggtagtgaa taccctactt caaagaacgg tgctgattca	1020
	actgctctag tcttcaggat ttagtacta ctgttttct gtttgaaca catggctgaa	1080
20	aataaatgtc tgcctttcga cctggcgct tagagaattt actaccacat ctcatcttta	1140
	gcatccaac gatgatttct gctgtcagaa tgaatgaatt gactaagagc aactcggtta	1200
	tttgagattg aattggttgt ttgtgattgt ttttgatttg tttttgctg tatgatcttt	1260
25	tgaggatttc gccatacaat gctgatacta gtcgtttgta tttccggta tatgtatttg	1320
	tgacgtatcg ttctgtagt ttgtaactaa tagaatgcat gtggtggtaa ctaatagaat	1380
30	gcatgttgta gtaacaaatg cacattgtag attctcgtgg attttcggg ttttcgttac	1440
	cagcacattg ccgattctgg tatgattttt gtcgtgttca ttgtttagt gcctttcttg	1500
	gctgccacta tttcattgag aatgtaggac gttgttcgat gcaaaagaac ttttccgac	1560
35	tagaatgcag gtggcaatct ggaatctctt attatgggag gaactactgt aattgggagg	1620
	ttttgattca gacaatctag taacagtcta gaagctactt tgcctttaaa tctcaatgac	1680
40	cttaaacgcc atgatggaga catttgaatc catgttttc aggtaaattg ggggcggctt	1740
	gacaaaatgg gagccggagg acgcatgtct gatccaacaa cgaaagacga acaaaagaag	1800
	aacccctcc aacgggtgcc ttacgcaaag cctccattca cactcggta catcaagaag	1860
45	gccattccac cacactgctt cgagagatcc gtcagccgtt cgttctcta tgcgtttac	1920
	gatctctca ttgttttct tctctactac attgcgactt ctacttcca tctgctgcca	1980
50	tcccatact gctacctagc ttggccattt tactgggctg tacaaggctg cgtttgacc	2040
	ggaatctggg tcattgcca tgaatgtggc caccatgcat tcagcgatta ccagtggctt	2100

WO 2004/044205

PCT/KR2003/002415

	gacgacacag ttggcctcat cctgcactct gccctgctcg tgccctattt ctcatggaaa	2160
	tacagccacc gccgccacca ctccaacact ggatcccttg agcgtgacga agtcttcgtc	2220
5	ccaaagccaa aatccagagt ctctgggtac tccaaatact tgaacaatcc acttggcaga	2280
	gtcatcacac ttgtggttac tcttactctc ggttggcctc tatacttgct gttaaatgc	2340
10	tctggcaggc cttacaaccg ttttgcagtc cactttgacc catatgggtcc aatatataat	2400
	gaccgtgaga gacttcaaat cttcatctcc gatgctggta taattgctgc tgtatgtgtg	2460
	ctttatcgtg ttgctttggt caaagggttg gcttggctgg tatgtgttta tggggtagcg	2520
15	ttactcattg tcaacgggtt ccttgttttg atcacattcc ttcagcacac tcacccttcg	2580
	ttgccgcact atgattcttc cgagtgggac tggctaaggg gagctcttgc aactgtcgac	2640
20	agagattatg ggggtgctaaa taagggttc cataacatca cagatacgca cgtgactcac	2700
	caccttttct caacgatgcc acattaccat gcaatggagg caactaaggc aatcaagccc	2760
	atactgggcc agtattatca gttgatgga accccgtttt acaaggcgat gtggagggag	2820
25	gcaaaggaat gtctgtatgt cgagccagac gagagtactc cagacaaggg tgtattctgg	2880
	tacaagaaca agttctgaag ccgaataaca tgtggttagt gaaaatggcg tcttcttatt	2940
30	ttgtctatg gagatggagg aacatcatca tgttttctt tttcttctat aagatgcgtc	3000
	ctttgttagt gtattctcig catgtaataa aataaacttc taccggaac cttgtctgtg	3060
	ctggtcggat tctagttctg caataaattg tcaagtttag tg	3102
35		
	<210> 2	
	<211> 383	
	<212> PRT	
40	<213> Sesamum indicum	
	<400> 2	
	Met Gly Ala Gly Gly Arg Met Ser Asp Pro Thr Thr Lys Asp Glu Gln	
	1 5 10 15	
45	Lys Lys Asn Pro Leu Gln Arg Val Pro Tyr Ala Lys Pro Pro Phe Thr	
	20 25 30	
	Leu Gly Asp Ile Lys Lys Ala Ile Pro Pro His Cys Phe Glu Arg Ser	
50	35 40 45	
	Val Ser Arg Ser Phe Ser Tyr Val Val Tyr Asp Leu Val Ile Val Phe	
	50 55 60	

WO 2004/044205

PCT/KR2003/002415

Leu Leu Tyr Tyr Ile Ala Thr Ser Tyr Phe His Leu Leu Pro Ser Pro  
 65 70 75 80

5 Tyr Cys Tyr Leu Ala Trp Pro Ile Tyr Trp Ala Val Gln Gly Cys Val  
 85 90 95

10 Cys Thr Gly Ile Trp Val Ile Ala His Glu Cys Gly His His Ala Phe  
 100 105 110

Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Leu His Ser  
 115 120 125

15 Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
 130 135 140

His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145 150 155 160

20 Pro Lys Ser Arg Val Ser Trp Tyr Ser Lys Tyr Leu Asn Asn Pro Leu  
 165 170 175

25 Gly Arg Val Ile Thr Leu Val Val Thr Leu Thr Leu Gly Trp Pro Leu  
 180 185 190

Tyr Leu Leu Phe Asn Val Ser Gly Arg Pro Tyr Asn Arg Phe Ala Cys  
 195 200 205

30 His Phe Asp Pro Tyr Gly Pro Ile Tyr Asn Asp Arg Glu Arg Leu Gln  
 210 215 220

Ile Phe Ile Ser Asp Ala Gly Ile Ile Ala Ala Val Cys Val Leu Tyr  
 225 230 235 240

35 Arg Val Ala Leu Val Lys Gly Leu Ala Trp Leu Val Cys Val Tyr Gly  
 245 250 255

40 Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Phe Leu  
 260 265 270

Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp Asp  
 275 280 285

45 Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Val Leu  
 290 295 300

Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Thr His His Leu  
 305 310 315 320

50 Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala Ile  
 325 330 335

WO 2004/044205

PCT/KR2003/002415

	Lys Pro Ile Leu Gly Gln Tyr Tyr Gln Phe Asp Gly Thr Pro Phe Tyr	
	340 345 350	
5	Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Leu Tyr Val Glu Pro Asp	
	355 360 365	
	Glu Ser Thr Pro Asp Lys Gly Val Phe Trp Tyr Lys Asn Lys Phe	
	370 375 380	
10		
	<210> 3	
	<211> 660	
15	<212> DNA	
	<213> Sesamum indicum	
	<220>	
	<221> promoter	
20	<222> (1)..(660)	
	<223> promoter of microsomal oleic acid desaturase coding gene	
	<400> 3	
25	catatgtgaa atgtaatgga aaatgcgaca agaattgcaa tagagaaaat ccaatttgca	60
	gagattacat gaaaagaatt tgtacaaata gcatatata gttaaatga aatgggacat	120
	gccacattat gtggaataaa aaagacaatt tgcttggaat taattataga ataatgtgt	180
30	tacatttaat atgtgattaa tcacttttt tgaattgtac atctatcaca tgacaagttc	240
	attatatttg acatataatt tgtttatgtc tagtcaagcc taattaaatt tctcgaaag	300
35	cacaaaattt tttgtccta accaggtttg aacaacaaa caaatcaca agcaggtgta	360
	tcgcacttgc gatgtgatcg gtcactttt cttaattgta catcattcac acgacaactg	420
	tattgtgctc caagttcaat tgagtgcggt tggagctata atttcttga acacacaatg	480
40	tggaatgtgc aactccatg tgggccaatg agcggatgac acgtggcggg caacttacct	540
	cgttacgttg aggcacatcat gaaaggggga tctcttgagg tggaggggtg ggggcggggg	600
45	ttgggggggg gccctctc agacaggtct atatttatga gacctcgtaa ggcagaacgc	660
		660



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/KR2003/002415

## A. CLASSIFICATION OF SUBJECT MATTER

IPC7 C12N 15/29

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7 C12N 15/29, C12N 15/00, C12N 9/00, C12N 9/06, C12N 15/82

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
Korean Patents and applications since 1975Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
NPS, PAJ, MEDLINE, KIPASS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT


Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2002/0061567 A (Y. Tom Tang, et al.) 23. 05. 2002 See the whole document	1-11
A	US 2002/0081691 A (Jennifer L. Hillman, et al.) 27. 06. 2002 See the whole document	1-11
A	US 2002/0045229 A (Ning Qin, et al.) 18. 04. 2002 See the whole document	1-11
A	US 2002/0166144 A (Allan Green, et al.) 07. 11. 2002 See the whole document	1-11
A	US 2002/0069426 A (Albert Boronat, et al.) 06. 06. 2002 See the whole document	1-11
A	US 2002/0059656 A (Qi Wang, et al.) 16. 05. 2002 See the whole document	1-11
A	KR 2001-0099065 A (POSTECH) 09. 11. 2001 See the whole document	1-11

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search  
04 FEBRUARY 2004 (04.02.2004)

Date of mailing of the international search report  
04 FEBRUARY 2004 (04.02.2004)

Name and mailing address of the ISA/KR  
 Korean Intellectual Property Office  
 920 Dunsan-dong, Seo-gu, Daejeon 302-701,  
 Republic of Korea  
 Facsimile No. 82-42-472-7140

Authorized officer  
 KWON, Oh Hee  
 Telephone No. 82-42-481-5597



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR2003/002415

Box No. 1 Nucleotide and/or amino acid sequence(s) (Continuation of Item 1.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:

a. type of material



a sequence listing



table(s) related to the sequence listing

b. format of material



in written format



in computer readable form

c. time of filing/furnishing



contained in the international application as filed



filed together with the international application in computer readable form



furnished subsequently to this Authority for the purposes of search

2. ☐ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR2003/002415

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2002/0061567 A	23. 05. 2002	EP 1268762 A CA 2399776 A AU 0133017 A	02. 01. 2003 09. 08. 2001 07. 08. 2001
US 2002/0081691 A	27. 06. 2002	US 6269192 A	05. 05. 2000
US 2002/0045229 A	18. 04. 2002	EP 1287134 A CA 2411069 A AU 0168212 A	05. 03. 2003 13. 12. 2001 17. 12. 2001
US 2002/0166144 A	07. 11. 2002	EP 0975765 A AU 6813598 A BR 9808676 A	02. 02. 2002 11. 11. 1998 11. 07. 2000
US 2002/0069426 A	06. 06. 2002	EP 1356033 A CA 2418436 A AU 0190522 A	29. 10. 2003 14. 02. 2002 18. 02. 2002
US 2002/0059656 A	16. 05. 2002	AU 0181464 A WO 016867 A	24. 09. 2001 20. 09. 2001
KR 2001-0099065 A	09. 11. 2001	None	